

N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED
IN THE INTEREST OF MAKING AVAILABLE AS MUCH
INFORMATION AS POSSIBLE

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

1057 F
81-10110
CR-164104

**THIRD QUARTERLY REPORT
FOR
EQUIVALENT SOURCE MODELING OF THE
MAIN FIELD USING MAGSAT DATA**

(E81-10110) EQUIVALENT SOURCE MODELING OF
THE MAIN FIELD USING MAGSAT DATA Quarterly
Report, 1 Jul. - 30 Sep. 1980 (Business and
Technological Systems, Inc.) 4 P
HC A02/MF A01

N81-22446

CSCL 08G G3/43 00110
Unclass

Contract NAS 5-26047

For the period
July 1, 1980 - September 30, 1980

submitted to

**NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
GODDARD SPACE FLIGHT CENTER
Greenbelt, Maryland 20771**

RECEIVED

JAN 9, 1981

SIS/902.6

M-022

TYPE II

by

**BUSINESS AND TECHNOLOGICAL SYSTEMS, INC.
Aerospace Building, Suite 440
10210 Greenbelt Road
Seabrook, Maryland 20801**

During this quarterly period the following modeling and software development work has been done.

- (1) A graphic software package has been developed to plot the dipole positions for a particular model on a world map. An arrow is drawn at each dipole in the direction of the horizontal magnetization vector, with length proportional to the horizontal magnitude. A contouring package represents the radial component of the magnetization vector. This software is in the final stage of verification.
- (2) Based on quiet MAGSAT data for November 5 and 6 (the MGST(6/80) data set), several dipole models defined in Table 1 have been derived. These models are being evaluated against one another and MGST(6/80) as to goodness of fit and stability of solution. The spatial power spectra computed from spherical harmonic expansions of the dipole models is being analyzed relative to crustal and core content. Figure 1 displays the power spectra for one of the dipole models and a spherical harmonic model based on MAGSAT data through degree and order 23.

Work planned for the next quarter includes further analysis of the derived models listed in Table 1 plus the derivation of an unconstrained $16^\circ \times 16^\circ$ dipole density model. The software will be enhanced to include time dependence (linear) in the dipole magnetization vector. This will facilitate the processing of available quiet MAGSAT data covering several months in time. It is anticipated that such a model will provide a useful tool for the reduction of MAGSAT data for anomaly studies.

Table 1

Model #	Dipole Density at core/mantle boundary	# dipoles	(constrained/ unconstrained) # degrees of freedom	Geocentric Dipole included	External field	RMS to Nov. 5, 6 MAGSAT data
1	32°x32°	42	126	NO	YES	33y
2	32°x32°	43	129	YES	YES	20y
3	21°x21°	92	276	NO	YES	7y
4	21°x21°	93	279	YES	YES	7y
5	16°x15°	162	162	NO	YES	7y

Figure 1. Power Spectra S_n of Magnetic Field Models

